

## University of Groningen

### Complementbindingsreactie volgens Witebsky, Klingenstein en Kuhn bij tuberculose

Beekhuis, Willem Henri

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1938

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Beekhuis, W. H. (1938). *Complementbindingsreactie volgens Witebsky, Klingenstein en Kuhn bij tuberculose*. [, Rijksuniversiteit Groningen]. [S.n.].

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## STELLINGEN.

### I.

Prophylactische bloed- of bloedseruminjecties van reconvalescenten of gezonde volwassenen verdienen in vele gevallen aanbeveling ter voorkoming of ter verzachting der mazelen.

### II.

De verschijnselen van de nephrose berusten op een glomerulus-aandoening.

### III.

Ook eiwitvrije stoffen kunnen als antigeen werken.

### IV.

De serologische antilichamen tegen hormonen berusten op de werking van de carbolgroep als antigeen.

### V.

In normale omstandigheden functioneert het foetale ademhalings-mechanisme.

### VI.

In de behandeling van de liesbreuken blijft voorloopig de operatieve behandeling verkieslijk boven de injectietherapie.

### VII.

Het is waarschijnlijk, dat bij de Morbus Paget mammae het carcinoom ontstaat in de groote melkgangen dicht bij de inmondingsplaats in den tepel, vanwaar het zich kan uitbreiden, zoowel in de huid als langs de melkgangen naar de diepte.

### VIII.

De behandeling der croupeuze pneumonie met antipneumococcenserum is een groote verbetering. Men bestede vooral aandacht aan het gebruik van konijnen-antipneumococcenserum.

## IX.

De lichaampjes van Gandy-Gamna vinden hun ontstaan in kleine bloedingen, onafhankelijk van eenige schimmel.

## X.

De behandeling van de fractura colli femoris intracapsularis met de metalen pen van Smith-Petersen of met die van Sven Johansson is verwerpelijk.

## XI.

Bij een patient met een peritonsillair absces verrichte men „tonsillectomie à tiède”.

## XII.

Bij de behandeling van endometriose heeft (röntgen) castratie niet altijd het gewenschte resultaat.

## XIII.

De beoordeeling van de psychische ontwikkeling in de eerste drie levensjaren is zonder grondige kennis der neurologie niet mogelijk.